

Exit gas data를 이용한 포도당 공급 제어 및 *Alcaligenes eutrophus* 에 의한 타피오카 당화액으로부터 PHB 생산

김범수, 장호남*

건양대 화학공학과, 한국과학기술원 화학공학과*

Control of Glucose Feeding Using Exit Gas Data and Its Application to the Production of PHB from Tapioca Hydrolysate by *Alcaligenes eutrophus*

Beom Soo Kim, Ho Nam Chang*

Department of Chemical Engineering, Konyang University,

Department of Chemical Engineering, KAIST*

SUMMARY

A method to estimate the glucose concentration in the culture broth using CER (CO₂ evolution rate) data from a mass spectrometer was developed. *Alcaligenes eutrophus* was cultivated to produce poly(3-hydroxybutyric acid) (PHB) from tapioca hydrolysate using this method. The *k* value (g glucose/mol CO₂) defined as the glucose consumption per CO₂ evolution decreased with culture time and was automatically changed using CER data. The glucose concentration in the culture broth could be controlled at 10 to 20 g/L. A final cell concentration of 106 g/L, PHB concentration of 61 g/L, and a P/X of 58 % were obtained after 59 h of cultivation.

INTRODUCTION

Poly(3-hydroxybutyric acid) (PHB) is a biodegradable thermoplastics which can be produced by many microorganisms as an environmentally harmless alternative to petrochemical derived plastics. The high production cost of PHB can be reduced by improving fermentation and separation processes and/or using a cheap carbon source.

Alcaligenes eutrophus has been most widely used organism for the production of PHB (Byrom, 1987; Doi, 1990). Recently, we found that maintaining glucose concentration at 10 to 20 g/L was important for efficient production of PHB by *A. eutrophus*, and obtained a high concentration of PHB (121 g/L) in a fed-batch culture with glucose concentration control using exit gas data from a mass spectrometer or using an on-line glucose analyzer (Kim et al., 1994).

When using CER data for glucose concentration control, however, it was needed to manually change the value of *k* (g glucose/mol CO₂) which represents the glucose consumption per CO₂ evolution. In this study, we developed a method to estimate the *k* value with time from a simple stoichiometry and an empirical method to automatically change the *k* value with time from the variables that could be monitored. Using this method, we carried out PHB production by *A. eutrophus* from tapioca hydrolysate.

MATERIALS AND METHODS

Organism and media

Alcaligenes eutrophus NCIMB 11599 was used in this study. The media for the seed culture was (per liter) glucose, 20 g; (NH₄)₂SO₄, 1 g; KH₂PO₄, 1.5 g; Na₂HPO₄·12H₂O, 9 g; MgSO₄·7H₂O, 0.2 g; trace element solution, 1 mL. The composition of the trace element solution was (per liter) FeSO₄·7H₂O, 10 g; ZnSO₄·7H₂O, 2.25 g; CuSO₄·5H₂O, 1 g; MnSO₄·4-5H₂O, 0.5 g; CaCl₂·2H₂O, 2 g; Na₂B₄O₇·10H₂O, 0.23 g; (NH₄)₆Mo₇O₂₄, 0.1 g; 35 % HCl, 10 mL.

For the fed-batch culture, the initial medium was (per liter) $(\text{NH}_4)_2\text{SO}_4$, 4 g; KH_2PO_4 , 13.3 g; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1.2 g; citric acid, 1.7 g; trace element solution, 10 mL. Tapioca hydrolysate containing 20 g/L glucose was added to the initial medium as a carbon source. The pH of the medium was adjusted to 6.8 with NaOH.

Preparation of tapioca hydrolysate

Tapioca hydrolysate was prepared from tapioca starch through liquefaction and saccharification processes with α -amylase and glucoamylase. The concentration of the total sugar after the saccharification process was about 250 g/L, and the composition was glucose, 89.9 %; maltose, 7.5 %; maltotriose, 0.3 %; etc., 2.3 %. The solution containing 500 g/L glucose was used as a feeding medium for the fed-batch culture, which was obtained by concentrating the tapioca hydrolysate using a rotary evaporator.

Cultivation and analytical procedures

The details on cultivation and analytical methods are described in our previous paper (Kim et al., 1994).

Control of glucose feeding using CER data from mass spectrometer

The feeding strategy is based on the assumption that both cell growth and PHB production are associated with CO_2 evolution. The glucose consumption rate r_s was expressed as


$$r_s = k \cdot \text{CER} \quad (1)$$

where k is a proportionality parameter which varies with culture time. The details on the determination of feeding rate are described in our previous paper (Kim et al., 1994).

RESULTS AND DISCUSSION

We found that high concentration of cell (124 g/L) and PHB (94 g/L) could be obtained when ammonia feeding was stopped at 55 g/L cell concentration from glucose as a carbon source by glucose concentration control using CER data (Kim et al., 1994). Since the value of k in equation (1) changed during the course of culture, we had to recalculate it manually at every sampling time. To solve this problem, we developed a method to change the k value automatically with culture time.

First, the theoretical k value was calculated by estimating the coefficients in the following simple stoichiometric equation.



It was assumed that glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) was consumed to produce cell ($\text{CH}_{1.666}\text{N}_{0.2}\text{O}_{0.27}$), PHB ($\text{C}_4\text{H}_6\text{O}_2$), and CO_2 . The chemical formula of the cell was taken from that of typical bacteria (Bailey and Ollis, 1986). Six coefficients (a , b , c , d , e , and f) were determined from balances on the four elements (C, H, O, and N) and two additional relationships provided by experimental determination of the yield of residual biomass from glucose ($Y_{R/S}$) and PHB yield from glucose ($Y_{P/S}$).

$$\text{C} : 6a = 1 + d + 4f \quad (3)$$

$$\text{H} : 12a + 3b = 1.666 + 2e + 6f \quad (4)$$

$$\text{O} : 6a + 2c = 0.27 + 2d + e + 2f \quad (5)$$

$$\text{N} : b = 0.2 \quad (6)$$

$$Y_{R/S} = \frac{20.7}{180a} \quad (7)$$

$$Y_{P/S} = \frac{86f}{180a} \quad (8)$$

The k value was calculated by the equation.

$$k = \frac{180\alpha}{d} \quad (9)$$

Fig. 1 shows the time course of k value. Empty squares and filled squares represent values determined from equation (9) and values actually used during the experiment, respectively. Both values continuously decreased with culture time even though they did not exactly coincide with each other.

Next, we investigated an empirical method to evaluate the k value more accurately. We found that the k values are associated with the ratio of CER at time t to total CO_2 evolution from time zero to t . Fig. 2 shows the relationship between k and the ratio of CER to total CO_2 evolution. The following empirical equation was obtained.

$$k(t) = \frac{134.1U(t)}{U(t) + 0.02065} \quad (10)$$

where

$$U(t) = \frac{CER_t}{\int_0^t CER dt} \quad (11)$$

The solid line in Fig. 2 represents the value calculated by equation (10) and agrees well to the values used during the experiments (filled squares). Also in Fig. 1, the estimated k values (solid line) from equation (10) correlated very well to the values used every sampling time during the experiment (filled squares). Therefore, glucose concentration could be controlled by automatically changing the k value with time using CER data.

This method was applied to the production of PHB from tapioca hydrolysate. The set point for the glucose concentration was 15 g/L. Fig. 3 shows the time courses of the concentrations of cell, PHB, glucose, and P/X. Nitrogen limitation was applied at the cell concentration of 67 g/L at 37.5 h. The final concentrations of cell and PHB obtained in 59 h were 106 and 61 g/L, respectively, resulting in a P/X of 58 %.

These values were lower than those of PHB (121 g/L) and P/X (75 %) when using purified glucose as a carbon source (Kim et al., 1994). It seems that some products including cyanide chemicals within the tapioca hydrolysate inhibited the cell metabolism. This phenomenon was also observed in the experiments on ethanol production by yeast from tapioca hydrolysate (Lee et al., 1994).

The glucose concentration in the culture broth was maintained at 10 to 20 g/L during the culture. This supports the adequacy of using CER data for the estimation of k value. This culture technique may be useful for the production of other metabolites when maintaining an optimum substrate concentration is important.

REFERENCES

- Bailey, J.E. and Ollis, D.F. (1986). *Biochemical Engineering Fundamentals*, 2nd ed., New York : McGraw-Hill.
- Byrom, D. (1987). *Trends Biotechnol.* 5, 246-250.
- Doi, Y. (1990). *Microbial Polyesters*, New York : VCH Publishers.
- Kim, B.S., Lee, S.C., Lee, S.Y., Chang, H.N., Chang, Y.K. and Woo, S.I. (1994). *Biotechnol. Bioeng.* 43, 892-898.
- Lee, W.G., Lee, Y.S., Chang, H.N. and Chang, Y.K. (1994). *Biotechnol. Techniques*, 8, 817-820.




Fig. 1 Time course of the k value.
 Solid line indicates the k value calculated from equation (10).
 (■) k value used during experiment
 (□) k value calculated from equation (9)

Fig. 2 Relationship between the k value and the ratio of CER to total CO₂ evolution at time t.
 Symbols are the same as in Fig. 1.




Fig. 3 Fed-batch culture of *A. eutrophus* from tapioca hydrolysate.
 (x) dry cell concentration (g/L); (■) PHB concentration (g/L)
 (□) P/X (%); (Δ) glucose concentration (g/L)